

KALAI ORIENTATIONS ON MATROID POLYTOPES

RAUL CORDOVIL

ABSTRACT. Let P a polytope and let $\mathcal{G}(P)$ be the graph of P . Following Gil Kalai, we say that an acyclic orientation $\mathcal{O}$ of $\mathcal{G}(P)$ is **good** if, for every non-empty face F of P , the induced graph $\mathcal{G}(F)$ has exactly one sink. Gil Kalai gave a simple way to tell a simple polytope from the good orientations of its graph. This article is a broader study of “good orientations” (of the graphs) on matroid polytopes.

Dedicated to Michel Las Vergnas on the occasion of his 65th birthday

1. INTRODUCTION

Let P be a simple polytope (see [10] for details on polytopes) and let $\mathcal{G}(P)$ be its graph (1-skeleton). M. Perles conjectured (see the reference [P] in [7]) and Blind and Mani [2] proved that the graph $\mathcal{G}(P)$ determines the lattice of faces of P . Kalai [7] gave a short and constructive prove of this result; see also [5, 6] for a discussion and refinement of Kalai’s technique. Kalai’s proof is based on an intrinsical characterization of the “good” orientations of $\mathcal{G}(P)$ between all the acyclic orientations. Following [7], we say that an acyclic orientation $\mathcal{O}$ of $\mathcal{G}(P)$ is **good** if for every non-empty face F of P the induced graph $\mathcal{G}(F)$ has exactly one **sink**, i.e., a vertex of $\mathcal{G}(F)$ with no lower adjacent vertices. Every linear ordering, $\{v_1 \prec \cdots \prec v_n\}$, of the vertex set V of $\mathcal{G}(\mathcal{M})$ induces an acyclic orientation $\mathcal{O}_\prec$ of the graph, where an edge is directed from its larger end-node to its smaller end-node. The linear ordering $\{v_1 \prec \cdots \prec v_n\}$ is called **good** if $\mathcal{O}_\prec$ is a good orientation. Each acyclic orientation of an arbitrary graph G is induced by some linear ordering of its vertices, see [9, Proposition 1.2]. Good orderings are in 1-1 correspondence with shelling orderings of the facets of the boundary ∂P^Δ of the dual polytope P^Δ (see Theorem 2.3 below for a matroidal generalization).

We say that an oriented matroid $\mathcal{M}$ is a **matroid polytope** if it is acyclic and all the elements of the ground set $E(\mathcal{M})$ are extreme points of $\mathcal{M}$. The **graph** $\mathcal{G}(\mathcal{M})$ of the matroid polytope $\mathcal{M}$ is the graph whose vertices [resp. edges] are the faces of rank 1 [resp. 2] of $\mathcal{M}$. In particular the vertex set of $\mathcal{G}(\mathcal{M})$ is the ground set $E(\mathcal{M})$. We say that the matroid polytope $\mathcal{M}$ is **simple** if every vertex of $\mathcal{G}(\mathcal{M})$ is incident with exactly $\text{rank}(\mathcal{M}) - 1$ edges.

The authors’ research was supported in part by FCT (Portugal) through program POCTI and the project “Algebraic Methods in Graph Theory” approved by program Pessoa 2005/6.

The proof of Gil Kalai [7] remains true if we replace “simple polytope” by “simple matroid polytope” so, the graph $\mathcal{G}(\mathcal{M})$ also encloses the face lattice of $\mathcal{M}$. If F is a non-empty face of $\mathcal{M}$ then F also is a matroid polytope (we are identifying F and the restriction of $\mathcal{M}$ to F). For more details on oriented matroid theory see Section 2 and [1].

2. K-ORDERINGS

Let $\mathcal{M}', \mathcal{M}''$ [resp. $\mathcal{M}$] denotes an acyclic oriented matroid [resp. matroid polytope] of rank r on ground set $E = \{e_1, \dots, e_n\}$. Let $\mathcal{O} = \mathcal{O}(\mathcal{M}')$ be the class of all the acyclic reorientations of $\mathcal{M}'$. For every linear ordering $E(\mathcal{M})_{\prec}$ let $\mathcal{O}_{\prec}$ denotes the orientation of $\mathcal{G}(\mathcal{M})$ where $\overrightarrow{uv}$ is a directed edge from u to v , if $v \prec u$. Let $\mathcal{G}_{\prec}(\mathcal{M}) := (\mathcal{G}(\mathcal{M}), \mathcal{O}_{\prec})$ be the corresponding digraph. If F is a face of $\mathcal{M}$ let $\mathcal{G}_{\prec}(F)$ be the induced directed subgraph on F . Note that an element e is the unique sink of $\mathcal{G}_{\prec}(\mathcal{M})$ if and only if for every element e' there is a directed path, $e' \rightsquigarrow e$, from e' to e .

Definition 2.1. *We say that the linear ordering $\{e_1 \prec e_2 \prec \dots \prec e_n\}$ of the ground set of a matroid polytope $\mathcal{M}$ is a **K-ordering** if, for every non-empty face F of $\mathcal{M}$, the directed subgraph $\mathcal{G}_{\prec}(F)$ has exactly one sink. In particular the element e_1 is the unique sink of the digraph $\mathcal{G}_{\prec}(\mathcal{M})$.*

From the “Topological Representation Theorem”, we know that there is a pure regular CW-complex of dimension $r - 1$, with the topology of a PL-sphere and encoding $\mathcal{O}$, see [1, Theorem 5.2.1]. We will denote this CW-complex by $\Delta(\mathcal{O})$, or by Δ for short, and called it the **CW-complex of acyclic reorientations of $\mathcal{M}'$** . To every cell $W \in \Delta$ we attach a “sign vector” $\sigma(W) \in \{0, +, -\}^{E(\mathcal{M})}$ (for precisions see [1]). We will identify W and $\sigma(W)$ and set $W_{(e_i)} = \sigma(W)_{(e_i)}$. The set

$$\text{supp}(W) = \{e_i : W_{(e_i)} \neq 0\}$$

is called the **support** of W . We say that the cell W' is a face of W if the following two conditions hold:

$$(2.1) \text{ supp}(W') \subset \text{supp}(W);$$

$$(2.2) \text{ For every } e_i \in E, \text{ if } e_i \in \text{supp}(W') \text{ then we have } W'_{(e_i)} = W_{(e_i)}.$$

A cell $P \in \Delta$ has dimension 0 (is a **vertex**) if, seen as a signed vector, P is a signed cocircuit of $\mathcal{M}$. We will see Δ as an “abstract” regular cell complex over the set of its vertices. Every cell $W \in \Delta$ is identified with the set of its vertices $V(W)$:

$$W \equiv V(W) := \{P : P \text{ is a vertex of } \Delta \text{ and } P \leq W\}.$$

The facets of Δ are called the **topes** and have support equal to $E(\mathcal{M})$. We use the letter T to denote a tope. Every tope $T \in \Delta$ is a PL-ball and its boundary ∂T a PL-sphere. Every tope T fixes one acyclic reorientation $\mathcal{M}'' \in \mathcal{O}(\mathcal{M}')$. To explicit this correspondence we write $T = T(\mathcal{M}'')$. (The opposite tope $-T$ fix the same oriented matroid $\mathcal{M}''$.) If a fixed acyclic oriented matroid $\mathcal{M}'$ is given we set $T(\mathcal{M}') = (+, +, \dots, +)$. Let $\mathcal{L}(\mathcal{M})$

[resp. $\mathcal{L}(T(\mathcal{M}))$] be the lattice of the faces (with both trivial faces) of $\mathcal{M}$ [resp. $T(\mathcal{M})$] ordered by inclusion. There is a canonical anti-isomorphism $\Xi : \mathcal{L}(\mathcal{M}) \rightarrow \mathcal{L}(T(\mathcal{M}))$ where, to every element X of rank s of $\mathcal{L}(\mathcal{M})$, $\Xi(X)$ denotes the element of rank $r - s$ of $\mathcal{L}(T(\mathcal{M}))$ determined by the following conditions:

$$\Xi(X)_{(e_\ell)} = \begin{cases} 0 & \text{if } e_\ell \leq X \\ + & \text{if } e_\ell \not\leq X. \end{cases}$$

The atoms of $\mathcal{L}(T(\mathcal{M}))$ are the image by Ξ of the co-atoms of $\mathcal{L}(\mathcal{M})$. The set of vertices and facets of CW-complex T , are respectively

$$\{\Xi(H) : H \text{ a facet of } \mathcal{M}\} \quad \text{and} \quad \{\Xi(e) : e \in E(\mathcal{M})\}.$$

If $\mathcal{M}$ is a simple matroid polytope then $T(\mathcal{M})$ is an (abstract) simplicial complex of dimension $\text{rank}(\mathcal{M}) - 1$. In particular, for every pair $\{W, W'\}$ of cells of $T(\mathcal{M})$, we have $V(W \wedge W') = V(W) \cap V(W')$. The following definition is a particular case of the standard one, see [3, 8, 10] for details. (The equivalence of Conditions (2.2.1) and (2.2.1') is left to the reader.)

Definition 2.2. Let Δ be the CW-complex of the acyclic reorientations of $\mathcal{M}$ and $T \in \Delta$ be the tope associated to $\mathcal{M}$. We say that the linear ordering $\{\Xi_1 := \Xi(e_1) \prec \Xi_2 \prec \dots \prec \Xi_n\}$ is a **shelling** of the PL-sphere ∂T and ∂T is **shellable** if one of the following equivalent conditions holds:

- (2.2.1) For every pair of co-atoms $\Xi_i \prec \Xi_j$ such that $\Xi_j \cap \Xi_i \neq \emptyset$, there is some facet $\Xi_\ell \prec \Xi_j$ such that $\Xi_\ell \cap \Xi_j$ is an abstract simplex of cardinality $r - 2$ of T and $\Xi_j \cap \Xi_i \subseteq \Xi_\ell \cap \Xi_j$;
- (2.2.1') For every pair of vertices $\{e_i, e_j\}$, $1 \leq i < j \leq n$, on a non-singular face F of $\mathcal{M}$, there is some $\ell < j$ such that $\vec{e_j e_\ell}$ is a directed edge of the digraph $\mathcal{G}_\prec(F)$.

Theorem 2.3. The following two statements are equivalent:

- (2.3.1) $\{e_1 \prec \dots \prec e_n\}$ is a K -ordering of $\mathcal{M}$;
- (2.3.2) $\{\Xi_1 = \Xi(e_1) \prec \Xi_2 \prec \dots \prec \Xi_n\}$ is a shelling of the PL-sphere $\partial T(\mathcal{M})$.

Proof. (2.3.1) $\implies$ (2.3.2). As $i < j$ we know that e_j is not a sink of $\mathcal{G}_\prec(F)$. So there is a directed edge $\vec{e_j e_\ell}$ of $\mathcal{G}_\prec(F)$ and (2.2.1') holds.

(2.3.2) $\implies$ (2.3.1). Let $\{e_{i_1} \prec \dots \prec e_{i_{n'}}\}$ be the induced ordering on a non-empty face F of $\mathcal{M}$. From Condition (2.2.1') we know that e_{i_1} is the unique sink of $\mathcal{G}_\prec(F)$ and (2.3.1) follows. $\square$

Let us recall that a linear ordering $\{e_1 \prec e_2 \prec \dots \prec e_n\}$ of the ground set $E(\mathcal{M}')$ is called a **shelling ordering** of $\mathcal{M}'$, if the orientation obtained from $\mathcal{M}'$ by changing the signs on the initial sets $E_i := \{e_1, e_2, \dots, e_i\}$, $i = 1, 2, \dots, n$, is also acyclic. Edmonds and Mandel proved that in this case, $\{\Xi_1 = \Xi(e_1) \prec \Xi_2 \prec \dots \prec \Xi_n\}$ is a shelling of the PL-sphere $\partial T(\mathcal{M})$, see [1, Proposition 4.3.1]. Note that if F is a face of a matroid polytope $\mathcal{M}$, every shelling ordering of $\mathcal{M}$ induces a shelling ordering on F . The following

result is a consequence of Theorem 2.3 and the above result of Edmonds and Mandel.

Corollary 2.4. *Every shelling ordering of a simple matroid polytope is also a K-ordering.* $\square$

Let $f_\ell(\mathcal{M})$ be the number of faces of rank $\ell + 1$ of $\mathcal{M}$, $-1 \leq \ell \leq r - 1$. By convention set $f_{-1}(\mathcal{M}) = f_{r-1}(\mathcal{M}) = 1$. By analogy with the definition of the **f**-vector and the **h**-vector of a polytope, we call the vector,

$$\mathbf{f}(\mathcal{M}) := (f_{-1}(\mathcal{M}), f_0(\mathcal{M}), f_1(\mathcal{M}), \dots, f_{r-2}(\mathcal{M}), f_{r-1}(\mathcal{M})),$$

the **f**-vector of the (simple) matroid polytope $\mathcal{M}$ and we call the vector,

$$\mathbf{h}^*(\mathcal{M}) := (h_0^*(\mathcal{M}), h_1^*(\mathcal{M}), \dots, h_{r-1}^*(\mathcal{M})),$$

determined by the formulas

$$(1) \quad h_\ell^*(\mathcal{M}) = \sum_{i=0}^{\ell} (-1)^{\ell-i} \binom{r-1-i}{\ell-i} f_{r-1-i}(\mathcal{M}), \quad \ell = 0, 1, \dots, r-1,$$

the **h**^{*}-vector of $\mathcal{M}$. Note that the **f**-vector also can be recovered from the **h**^{*}-vector:

$$(2) \quad f_\ell(\mathcal{M}) = \sum_{i=0}^{r-1-\ell} \binom{r-1-i}{\ell} h_i^*(\mathcal{M}), \quad \ell = 0, 1, \dots, r-1.$$

(See [10, Section 8.3] for a good survey of **h**-vectors of simplicial polytopes and Dehn-Sommerville Equations.) From Euler-Poincaré formula (see [1, Corollary 4.6.11]) we know that

$$(3) \quad \sum_{i=-1}^{r-1} (-1)^i f_i(\mathcal{M}) = 0.$$

The graph $\mathcal{G}(\mathcal{M})$ is regular of degree $r-1$. Fix a K-ordering $\{e_1 \prec \dots \prec e_n\}$ of $\mathcal{M}$. Let $d^+(e)$ [resp. $d^-(e)$] denotes the outdegree [resp. indegree] of $e \in E$. Set

$$\mathbf{d}_\ell^+(\mathcal{M}) := |\{e : e \in E, d^+(e) = \ell\}|,$$

$$\mathbf{d}_\ell^-(\mathcal{M}) := |\{e : e \in E, d^-(e) = \ell\}|.$$

We clearly have that $\mathbf{d}_\ell^+(\mathcal{M}) = \mathbf{d}_{r-1-\ell}^-(\mathcal{M})$.

Theorem 2.5. *The integers $\mathbf{d}_\ell^+(\mathcal{M})$, $\ell = 0, 1, \dots, r-1$, are invariant of the matroid polytope $\mathcal{M}$ (i.e., are independent of the K-ordering) and they are determined by the equalities:*

$$(4) \quad h_\ell^*(\mathcal{M}) = \mathbf{d}_\ell^+(\mathcal{M}), \quad \ell = 0, 1, \dots, r-1.$$

Proof. Note that

$$\begin{aligned} \mathbf{f}_\ell(\mathcal{M}) &= \sum_{j=\ell}^{r-1} \binom{j}{\ell} \mathbf{d}_j^-(\mathcal{M}) \\ &= \sum_{i=0}^{r-1-\ell} \binom{r-1-i}{\ell} \mathbf{d}_{r-1-i}^-(\mathcal{M}) \\ &= \sum_{i=0}^{r-1-\ell} \binom{r-1-i}{\ell} \mathbf{d}_i^+(\mathcal{M}). \end{aligned}$$

From Equation (2) we conclude that $\mathbf{d}_\ell^+(\mathcal{M}) = h_\ell^*(\mathcal{M})$. $\square$

Corollary 2.6. *Let $E_{\prec} := \{e_1 \prec e_2 \prec \cdots \prec e_n\}$ be a K -ordering of $\mathcal{M}$. Then the reverse ordering $E_{\prec^*} := \{e_n \prec^* e_{n-1} \prec^* \cdots \prec^* e_1\}$ is also a K -ordering of $\mathcal{M}$.*

Proof. It is necessary to prove that, for every non-empty face F of $\mathcal{M}$, the digraph $\mathcal{G}_{\prec^*}(F)$ has exactly one sink, i.e., $\mathbf{d}_{\text{rank}(F)-1}^+(F) = 1$. From the equalities (4) we know $\mathbf{d}_{r-1}^+(\mathcal{M}) = h_{r-1}^*(\mathcal{M})$. From Euler-Poincaré formula (3) we conclude that

$$(5) \quad h_{r-1}^*(\mathcal{M}) = \sum_{i=0}^{r-1} (-1)^{r-1-i} f_{r-1-i}(\mathcal{M}) =$$

$$(6) \quad = \sum_{j=0}^{r-1} (-1)^j f_j(\mathcal{M}) = f_{-1}(\mathcal{M}) = 1.$$

So $\mathcal{G}_{\prec^*}(M)$ has exactly one sink. As every face F of $\mathcal{M}$ is a simple matroid polytope the result follows. $\square$

3. THE CUBE

Let $C^d := \{x \in \mathbb{R}^d : 0 \leq x_\ell \leq 1, \ell = 1, \dots, d\}$ be the d -dimensional cube. As the polar of the cube C^d is the d -dimensional crosspolytope (a simplicial polytope), it results from Theorem 2.5 above that:

$$(7) \quad \mathbf{h}^*(C^d) = \left(\binom{d}{0}, \binom{d}{1}, \dots, \binom{d}{d-1}, \binom{d}{d} \right),$$

i.e., there are exactly $\binom{d}{\ell}$, $0 \leq \ell \leq d$, vertices of $\mathcal{G}(C^d)$ such that $\mathbf{d}^-(e) = d - \ell$.

Let $B := \{0, 1\}^d$ be the set of the vertices of the cube C^d and for every $b_i \in B$, $1 \leq i \leq 2^d$, set $e_i = (b_i, 1) \in \mathbb{R}^{d+1}$. The rank $d+1$ **cube matroid polytope**, $\mathcal{C}^d$, is the oriented matroid determined by the linear dependencies of vectors of $E := \{e_i : b_i \in B\}$. The following theorem is closely related to the results presented here. (We present here a slightly different version of the original result.)

Theorem 3.1. [4, Proposition 3.2] *Let $\mathcal{C}^d$ be the cube matroid polytope of rank at least three. Let $\{e_1 \prec e_2 \prec \cdots \prec e_{2d}\}$ be a linear ordering of $E(\mathcal{C}^d)$. Then the following two conditions are equivalent:*

- (3.1.1) *$\{e_1 \prec e_2 \prec \cdots \prec e_{2d}\}$ is a K-ordering of $\mathcal{C}^d$;*
- (3.1.2) *For every rank three face F of $\mathcal{C}^d$, the digraph $\mathcal{G}_{\prec}(F)$ has an unique sink.* □

In the rank three cube matroid polytope $\mathcal{C}^2$, the “K-ordering” and the “shelling orderings” coincide. This result suggest the following problem:

Open Problems 3.2. *Is there a simple characterisation of shelling orderings of the cube matroid polytope $\mathcal{C}^d$? Are there K-orderings of the cube matroid polytope that are not shelling orderings?*

REFERENCES

- [1] Björner, A.; Las Vergnas, M.; Sturmfels, B.; White, N.; Ziegler, G.M.: Oriented Matroids. Second edition. Encyclopedia Math. Appl., **46**, Cambridge University Press, Cambridge, 1999.
- [2] Blind, Roswitha; Mani-Levitska, Peter: Puzzles and polytope isomorphisms. *Aequationes Math.* **34** (1987), 287–297.
- [3] Danaraj, Gopal; Klee, Victor: Which spheres are shellable?. *Ann. Discrete Math.* **2** (1978), 33–52.
- [4] Hammer, P. L.; Simeone, B.; Liebling, Th. M.; de Werra, D.: From linear separability to unimodality: a hierarchy of pseudo-Boolean functions. *SIAM J. Discrete Math.* **1** (1988), 174–184.
- [5] Joswig, Michael; Kaibel, Volker; Körner, Friederike On the k -systems of a simple polytope. *Israel J. Math.* **129** (2002), 109–117.
- [6] Kaibel, Volker: Reconstructing a simple polytope from its graph. *Lect. Notes Comput. Sci.* **2570**, Springer, Berlin, 2003, 105–118.
- [7] Kalai, Gil: A simple way to tell a simple polytope from its graph. *J. Combin. Theory Ser. A* **49** (1988), no. 2, 381–383.
- [8] Kleinschmidt, Peter; Smilansky, Zeev New results for simplicial spherical polytopes. *DIMACS, Ser. Discret. Math. Theor. Comput. Sci.* **6** (1991), 187–197.
- [9] Mihalisin, J.; Klee, V.: Convex and linear orientations of polytopal graphs. *Discrete Comput. Geom.* **24** (2000), no. 2-3, 421–435.
- [10] Ziegler, G. M.: Lectures on Polytopes. *Graduate Texts in Mathematics* **152**, Springer, Heidelberg, 1995.

DEPARTAMENTO DE MATEMÁTICA,
INSTITUTO SUPERIOR TÉCNICO,
AV. ROVISCO PAIS, 1049-001 LISBOA (PORTUGAL).
E-mail address: CORDOVIL@MATH.IST.UTL.PT